Missouri Botanical Garden

Peter H. Raven Library

George Engelmann Botanical Notebooks



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Rio Ku Sept 1869

Rhopalis,

in arboribus pendula, Prov. San Paulo, Bras.

Burchell No 3691

"trichotomus in ramos persecutis, in
lemniscis, dichotomus; bracteae globosae
albidae purpureo-rutilae non deciduae
(quod in Rh. Casupia!) diam. 2. Chm."

2331

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Isisoacty

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

H. Martius Aug 1869

368 (" Rhipsalis Cassyta, F." Forma robustior, articulis elongatis, floribus sublaeviusculis " Lagoa 1857)
J. A. Lindberg, Sept 1854. Minas Geraës, in silvis
flowers large, stem slender thin — in the order
Jan — allied to it — v to Jundiaï?

1479 Willdschlaegel, 14 arboris — Pernambuco —
very slender, with very small flowers,
name 1857 by me Rh. pauciflora
= H. Martii — Surinam, Paramaibo
branches articulate, or attenuated (3—8)
from near the ends of the older ones

364 Rhipsalis San Paolo Brasil = epiphyte — J. A. Lindberg.
Apr. 1854 — flowers extremely small —
perhaps same as N. 366. of the same collector —
(see last page) — 3—5 bracteolate from end
of branches.

367 coll by same = Nov 1854, Minas Geraës, Brazil
= similar, but still more slender, branches
bracteately divided — Ovaries = right angles,
no fl = fr.

1478 Rh. thomboidea ! Willdschlaegel Paramaibo,
Surinam — " in arboribus ; fl. pauci albi "
no fl. on specimen

2332

611 Rhipsalis Minas Geraës in Brasil
arborum, J. A. Lindberg Oct 1854
joint like Dissocactus — 25—35 cent long 1½—2½
wide — flat. — Same among Salzmann's

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Fort Dyck Jan. 1857

2333

Rhipsalis Cassyta is now covered with white berries — it is twelfth

Rh. ramulosa, unripe fruit, it is narrowly winged

Rh. rhombea, Swartziana, rispata are broad winged, yellow or yellowish flowers now open

Rh. pachyptera is Hooker, else similar, now in bloom

Rh. pentaptera very curious arrangement.

Rh. micrantha similar, 5 angled. but much more slender

Missouri Botanical Garden
George Engelmann Papers

Cauteac Hort. Venet.

Jan 1858

Physalis fraxifolia is more pendulous;
Ph. flavescens is similar, but more
erect; both thick. —

2334

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Rhipsalis

Frankfurt Febr. 1857

2335



Rhipsalis Cassyta Hort Dyck

seed 0.6 lin long, subterete, linear rounded on back, subarcuate in front, hilum subbasal, oblong linear; testa brittle, dark chestnut brown shining, finely striolate

Embryo subterete straight terete, (rectus) albumen none subclavate; indicated by a slightly thickened brownish membr. cotyledons foliaceous, erect, (contrary) parallel acutish, abot ¼ the length of the whole embryo. —

Rh. ramulosa Hort Dyck

seed 0.6 lin long, obovate, compressed not keeled; hilum subbasal, oblong linear; testa chestnutbrown, shining brittle, very slightly reticulose;



fruit white transparent, globose,
fl. persistent.

Rh. crispata (Hort Jacobi)

seed 0.45–0.5 obliquely obovate, compressed; chestnutbrown, shining almost smooth; hilum, obliquely basilar linear-oblong; seed cavity anteriorly.

Embryo 0.4 compressed; erect. albumen obovate compressed, cotyledons contrarious, short erect, about ¼ of the whole Embryo, laterally compressed.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Rhipsalis Jan 1860

fruit or rather ovary with as many cells as style has lobes ! No!

Ja'ckn. Struct t. 136 tab 28 (1788) says that the fruit has the seeds attached to a central axis

Hooker (ubi?) fide DC. Rev. p. 78 repeats this — DC. Prod. III. p 475 repeats it.

DC. Rev. (1829) p. 79 (see also p. 18) for the first time states that the figure of the Flor. Mex. and Gay Moquin & Serre, figures the fruit as tricoccous & he copies the figure pl. 21. — but is doubtful whether there are 3 or a single cell, — he adheres to the opinion of the central attachment of the seeds 1 years later Pr Salm. — Otto Gord Zeit 1836 p. 146 and Meissn Gen. p. 185 quoted, after careful examination that the fruit of Rhipsalis is identical with the fruit of all other Cactaceae

2336

Both Gaertner and the Mexican flora are right to some extent

The ovary has as many cells as the style has lobes ! with a thick fleshy axis

But the ovules are not attached to the axis but to the parietal placenta which reaches like an incomplete dissepiment into the cell, dividing it almost completely into 2 halfs

The ovula hang from the inner edge of the placenta, which is perfectly free from the axis, into the semicells their base being attached (the reverse of other Cacti with short funiculi) directed to the orifice of the fruit, their micropyle to the outer

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

9

flowers often larger than the plant itself;
these forms extend towards into New Mexico
and into the northern provinces of Mexico
but appear to be unknown further
south, — where the species of the family
are most numerously developed. —

Other genera of this peculiarly
american family have not been
found in Texas. J.

There are some other plants of
different families characteristic of
that flora — as the Ruellia Texanum
which with Pegamum (A. G. Mss)
of the Rio Grande Mexican and south
representatives of the Asclepiadeae
America. — So also the remarkable
Hermannia Texana, those congeners
are so common South Africa;

A. As we may expect the flora
of Southwestern Texas partakes to some
extent of the characters of the
vegetation of the adjoining districts. —
Thus. We find there the Oenotheras
and Gaurae of the Buffalo Grass region of the
Plains & the now well known compositae Plant, Silphium laciniatum is also
C. The Western mountain region see page 10.
sends down to the calcareous plateaus B.
a number of grasses & some Agrostideae, etc
a goodly number Tribes
From Texas the Plateau of Mexico and
particularly its northern slopes we
find in our flora the Boliviaceae
(some of the different species to be sure)
some Malpighiaceae, some Zygophyllaceae
besides the Mimoseae & Cactaceae
to be already mentioned; Mostly genera or
families not found further North and east.

D.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Rhipsalis 2 Jan 1860 N. 2

2337

There is some resemblance to the Cereeæ
fruit, but in this the placentæ are curled
spirally, forming part of the dissepiment there
& there: & then
and also the central axis



In Rhipsalis it would seem that the central axis
& the dissepiment is formed by the protruding
and centrally uniting midrib of the carpels.

These investigations were made Jan 19th on Rh. pachyptera

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

Turpin Obs. sur la fam. des Cactées (1830) says
pag. 62 that the ovary of Rhipsalis & 3 locular, the
ovules attached to the internal angles of the cells
pl. 3 fig 5 he figureth the ovary thus:



Pfeiffer Enum. 1837. says pag: 136: "Delineatio
horum in Turpini obs. omnino falsa".
Salm Dyck. 1850. p. 230: in generibus omnibus Cact. semina parietalia (pointed to Rhipsalis

Jan 25. A new examination of the fruit shows that I was
mistaken! There are curtains by the placentæ
protruding into the cavity of the ovarian cavity
with removed ovules, pretty nearly as Turpin
has it — these forming an incompletely 3—5—
locular ovary — but no other dissepiments
are there — The fruit is therefore then
like the Pepo fruit, but
placentæ not curled up as in
that fruit. —





placentæ & ovules
seen from the inside — center of ovary

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

so different from that of Texas in the same
latitude; thus the western slope of
the Alleghanies towards the immediate valley
of the Mississippi, the northern and the
southern part of that valley itself, the
latter including northeastern Texas; then again
the region of the western plains and
deserts and finally the peculiar
flora of Southwestern Texas or the flora
of the Rio Grande valley.

F see next page
& page 5 & 6

The peculiar character of the vegetation
of this region consists in the diminution of
trees into shrubs; they do not disappear
entirely as in the western plains and
deserts which extend into northwestern
Texas, nor are generally the same species which
elsewhere grow up to be large trees, greatly
reduced in size, but shrubby species
peculiar to this region represent the
larger trees of the same or analogous
genera of the more northern parts of the country.
The stately Walnuts of your forests
are reduced there to the low Juglans nana,
which bears nuts of the size of a marketball. In place of
the Mulberry of your river bottoms we find
there a shrub, with leaves of 1/4 the size
the Morus parvifolia. The fine Hackberry,
Celtis occidentalis, one of the largest
trees of the fertile lands of the western
country is there represented by a
analogous allied spiny shrub of a nearly
allied genus which not being able
to find a description of any work at
my disposal I have named Acanthoceltis
assumed to

2338

Gaerta de fruct.

I p. 137

Rhipsalei . . .

. . . . Cacta i.afera
uni.loculari.s pellucida, semina
d.uodena centralia

Cacta i.afera flori.
_ ued'i.mento coronata, sphaerica
nuca pellucida uni.loculari.s,
Polpa aqueosa, seminibus nudique
circumfusa

Recep. nullum, praeter vasa nutritia
semina ad axem fructus alligata.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS